arXiv:1608.04131v1 [math.DG] 14 Aug 2016

NULL SECTIONAL CURVATURES OF WARPED PRODUCTS

BENGİ R. YAVUZ, BÜLENT ÜNAL, AND FERNANDO DOBARRO

Abstract. In this paper, we investigate the null (light-like) sectional curvatures of Lorentzian warped product manifolds. We derive the formulas for the null sectional curvature of many well-known warped product space-time models such as multiply generalized Robertson-Walker space-times, generalized Kasner space-times and standard static space-times.

Contents

1. Introduction 1
2. Preliminaries 3
3. Null Sectional Curvature 6
3.1. Null Sectional Curvature of a multiply GRW space-time 6
3.2. Null Sectional Curvature of a GRW space-time 8
3.3. Null Sectional Curvature of a Generalized Kasner Space-time 9
4. Four-dimensional Space-time Models 10
4.1. Type I: Null Sectional Curvature of a Kasner Space-time with fiber of dimension (3) 10
4.2. Type II: Null Sectional Curvature of a Kasner Space-time with fiber of dimension $(1, 2)$ 10
4.3. Type III: Null Sectional Curvature of a Kasner Space-time with the fiber of dimension $(1, 1, 1)$ 11
5. Null sectional curvature of a SSS-T 12
Appendix A. Null Sectional Curvature of MGRW space-time 13
References 16

Date: July 26, 2016.
2010 *Mathematics Subject Classification.* Primary 53C25; Secondary 53C80.
Key words and phrases. Warped products, null sectional curvature, warped product space-times.

1. Introduction

Warped product manifolds were first introduced to the literature by R. Bishop and B. O'Neill in [7] to construct complete Riemannian manifolds with negative sectional curvature everywhere. After that Beem, Ehrlich and Powell established that a wide class of well known exact solutions of the Einstein's field equations can be expressed as a Lorentzian warped product in [4, 5]. Two important examples of warped products are the generalized Robertson-Walker space-times (GRW, for short) and the standard static space-times (SSS-T, for short). The former are obviously a generalization of Robertson-Walker space-times and the latter a generalization of the Einstein static universe. In addition to these space-times, we can list generalized Kasner and Reissner-Nordström space-time models.

In the current paper, we focus on the null (light-like) sectional curvatures of Lorenztian warped products. The concept of null sectional curvature were first defined by S. G. Harris in [14] to study sectional curvatures of null plane sections of Lorentzian manifolds. In [15], it is proved that Robertson-Walker metrics can be locally characterized as those for which the null sectional curvature denoted by K_N at each point is a constant for all the null planes at that point.

In [24], J. P. Palomo proved the smoothness of the null sectional curvature. Moreover, in a not necessarily complete manifold, if the null sectional curvature is not zero for any degenerate planes at a point p, then there is at most one local GRW structure in a neighborhood of p (see [12]). Moreover he further showed that for an $n(\geq 4)$-dimensional manifold the $U-$ normalized null sectional curvature function is constant on every pencil of degenerate planes belonging to a certain null direction, if and only if, the Lorentzian manifold is conformally flat.

In [15, 19], it is shown that if (M, g) is a time orientable Lorentzian manifold with $dim(M) \geq 3$, and U is a globally defined unitary time-like vector field, then the $U-$ normalized null sectional curvature is isotropic, i.e, it is only a point function. Moreover,

$$K^U(p, \Pi) = K^U(p)$$

for all null planes Π if and only if the curvature tensor satisfies:
(a) $R(X, Y)Z = k(X \wedge_g Y)Z$, for any $X, Y, Z \in U^\perp$
(b) $R(X, U)U = \mu X$, for any with $\kappa, \mu \colon M \to \mathbb{R}$ and $K^U = \kappa + \mu$.

If the $U-$ normalized null sectional curvature K^U is isotropic (it is said to be spatially constant) if it is constant on the space of orthogonal to the chosen time-like vector field U, i.e, $K^U = 0$ for every $X \in U^\perp$ (see [19]).

Let (M, g) be an $n \geq 4$- dimensional Lorentzian manifold and let U be a unitary time-like vector field on M. Suppose that the U-normalized null sectional curvature is non-zero, isotropic and spatially constant. Then g is locally a Friedmann-Lemaitre-Robertson-Walker metric (see [15, 19]).

In [12], to characterize global decomposition of a manifold as a generalized Robertson-Walker space-time, the null sectional curvature is applied.

The null sectional curvature has been used in the study of conjugate points along null geodesics (see [13, 14, 25]).

We recall that a *warped product* can be defined as follows [5, 23]. Let (B, g_B) and (F, g_F) be pseudo-Riemannian manifolds and also let $b\colon B \to (0, +\infty)$ be a smooth function. Then the (singly) warped product, $B \times_b F$ is the product manifold $B \times F$ furnished with the metric tensor $g = g_B \oplus b^2 g_F$, more precisely

$$g = \pi^*(g_B) \oplus (b \circ \pi)^2 \sigma^*(g_F),$$

where $\pi\colon B \times F \to B$ and $\sigma\colon B \times F \to F$ are the usual projection maps and * denotes the pull-back operator on tensors.

A *standard static space-time* can be considered as a Lorentzian warped product where the warping function is defined on a Riemannian manifold called the fiber and acting on the negative definite metric on an open interval of real numbers, called the base. More precisely, a SSS-T, ${}_f(a, b) \times F$ is a Lorentzian warped product furnished with the metric $g = -f^2 \mathrm{d}t^2 \oplus g_F$, where (F, g_F) is a Riemannian manifold, $f\colon F \to (0, +\infty)$ is smooth, and $-\infty \leq a < b \leq +\infty$. In [23], it was shown that any static space-time is locally isometric to a SSS-T.

Standard static space-times have been previously studied by many authors. Kobayashi and Obata [18] stated the geodesic equation for this class of space-times and the causal structure and geodesic completeness was considered in [3], where sufficient conditions on the warping function for nonspacelike geodesic completeness of the SSS-T was obtained (see also [26]). In [2], conditions are found which guarantee that SSS-Ts either satisfy or else fail to satisfy certain curvature conditions from general relativity. The existence of geodesics in SSS-Ts have been studied by several authors. Sánchez [27] gives a good overview of geodesic connectedness in semi-Riemannian manifolds, including a discussion for SSS-Ts.

Two of the most famous examples of SSS-Ts are Minkowski space-times and the Einstein static universe [5, 17] which is $\mathbb{R} \times \mathbb{S}^3$ equipped with the metric

$$g = -\mathrm{d}t^2 + (\mathrm{d}r^2 + \sin^2 r \mathrm{d}\theta^2 + \sin^2 r \sin^2 \theta \mathrm{d}\phi^2)$$

where $\mathbb{S}^3$ is the usual 3-dimensional Euclidean sphere and the warping function $f \equiv 1$. Another well-known example is the universal covering space of the anti-de Sitter space-time, a SSS-T of the form ${}_f\mathbb{R} \times \mathbb{H}^3$ where $\mathbb{H}^3$ is the 3-dimensional hyperbolic space with constant negative sectional curvature and the warping function $f\colon \mathbb{H}^3 \to (0, +\infty)$ defined as $f(r, \theta, \phi) = \cosh r$ [5, 17]. Finally, we can also mention the Exterior Schwarzschild space-time [5, 17], a SSS-T of the form ${}_f\mathbb{R} \times (2m, +\infty) \times \mathbb{S}^2$, where $\mathbb{S}^2$ is the 2-dimensional Euclidean sphere, the warping function $f\colon (2m, +\infty) \times \mathbb{S}^2 \to (0, +\infty)$ is given by $f(r, \theta, \phi) = \sqrt{1 - 2m/r}$ and the

line element on $(2m, +\infty) \times \mathbb{S}^2$ is

$$\mathrm{d}s^2 = \big(1 - \frac{2m}{r}\big)^{-1}\mathrm{d}r^2 + r^2(\mathrm{d}\theta^2 + \sin^2\theta \mathrm{d}\phi^2).$$

2. Preliminaries

In this section, we give the formal definitions of several types of warped product space-time models such as GRWs, SSS-Ts and multiply generalized Robertson-Walker space-times (MGRW, for short) and their related geometric structure formulas (see [5, 23]).

Throughout the article I will denote an open real interval of the form $I = (t_1, t_2)$, where $-\infty \leq t_1 < t_2 \leq +\infty$.

Definition 1. *Let (F, g_F) be an s-dimensional Riemannian manifold and $b\colon I \to (0, +\infty)$ be a smooth function. Then the $n(= 1 + s)$-dimensional product manifold $I \times F$ furnished with the metric tensor $g = -\mathrm{d}t^2 \oplus b^2 g_F$ is called a generalized Robertson-Walker space-time and is denoted by $I \times_b F$, where $\mathrm{d}t^2$ is the Euclidean metric tensor on I.*

Definition 2. *Let (F, g_F) be an s-dimensional Riemannian manifold and $f\colon F \to (0, +\infty)$ be a smooth function. Then the $n(= 1 + s)$-dimensional product manifold $I \times F$ furnished with the metric tensor $g = -f^2\mathrm{d}t^2 \oplus g_F$, where $\mathrm{d}t^2$ is the Euclidean metric tensor on I, is called a standard static space-time and is denoted by ${}_f I \times F$.*

Definition 3. *Let (B, g_B) and (F_i, g_{F_i}) be pseudo-Riemannian manifolds and also let $b_i\colon B \to (0, \infty)$ be smooth functions for any $i \in \{1, 2, \cdots, m\}$. The product manifold $M = B \times F_1 \times F_2 \times \cdots \times F_m$ furnished with the metric tensor $g = g_B \oplus b_1^2 g_{F_1} \oplus b_2^2 g_{F_2} \oplus \cdots \oplus b_m^2 g_{F_m}$ is called a multiply warped product and is denoted as*

$$B \times {}_{b_1} F_1 \times \cdots \times {}_{b_m} F_m.$$

More precisely

$$g = \pi^*(g_B) \oplus (b_1 \circ \pi)^2 \sigma_1^*(g_{F_1}) \oplus \cdots \oplus (b_m \circ \pi)^2 \sigma_m^*(g_{F_m}), \tag{2.1}$$

where the π and σ_i are the usual projection maps and * denotes the pull-back operator on tensors. (B, g_B), $b_i\colon B \to (0, \infty)$ and (F_i, g_{F_i}) are called base, i-fiber and i-warping function of the multiply warped product, respectively.

- If $m = 1$, then we obtain a *singly warped product*.
- If all $b_i \equiv 1$, then we have a (trivial) *product manifold*.
- If (B, g_B) and all the (F_i, g_{F_i}) are *Riemannian* manifolds, then (M, g) is also a *Riemannian* manifold.
- The multiply warped product (M, g) is a *Lorentzian multiply warped product* if (F_i, g_{F_i}) are all *Riemannian* for any $i \in \{1, 2, \cdots, m\}$ and either (B, g_B) is *Lorentzian* or else (B, g_B) is a one-dimensional manifold with a *negative definite* metric $-\mathrm{d}t^2$.

- If B is an open interval I equipped with the negative definite metric $g_B = -\mathrm{d}t^2$ and all the (F_i, g_{F_i}) manifolds are Riemannian, then the Lorentzian multiply warped product (M, g) is called a *multiply generalized Robertson-Walker space-time*. In particular, a MGRW is called a *generalized Reissner-Nordström space-time* when $m = 2$.

Throughout the paper the fiber(s) (F, g_F) of a warped product space-time model is always assumed to be connected. We denote the set of lifts of vector fields on B and F to $B \times F$ by $\mathfrak{L}(B)$ and $\mathfrak{L}(F)$, respectively and use the same notation for a vector field and for its lift (see page 205 of [23]).
From now on, we follow the convention applied in [5] (note the difference with [23]) for the definition and sign of the Riemann curvature tensor R, namely. For any n-dimensional pseudo-Riemannian manifold (N, h),

$$R(X, Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X,Y]} Z,$$

where ∇ is the h-Levi-Civita connection and X, Y, Z are vector fields on N. Besides, Ric will denote the Ricci tensor (see [23]).
Furthermore, we will apply the sign convention for the Laplacian in [23], i.e, $\Delta = \mathrm{tr}(\mathrm{H}) = \mathrm{div}\,\mathrm{grad}$, where tr denotes the h-trace, H the Hessian tensor respect to the Levi-Civita connection, div the h-divergence and grad the h-gradient (see page 85 of [23]).
Eventually, we will put a sub- or super-index in the corresponding operator indicating the manifold on which it is acting, for instance ∇^B if the connection is that on the manifold B.

The following basic formulas about the geometry of multiply warped product can be found in [10]

Proposition 1. *Let $M = B \times_{b_1} F_1 \times \cdots \times_{b_m} F_m$ be a pseudo-Riemannian multiply warped product with metric $g = g_B \oplus b_1^2 g_{F_1} \oplus \cdots \oplus b_m^2 g_{F_m}$ also let $X, Y \in \mathfrak{L}(B)$ and $V \in \mathfrak{L}(F_i)$, $W \in \mathfrak{L}(F_j)$. Then*

(1) $\nabla_X Y = \nabla^B_X Y$

(2) $\nabla_X V = \nabla_V X = \dfrac{X(b_i)}{b_i} V$

(3) $\nabla_V W = \begin{cases} 0 & \text{if } i \neq j \ , \\ \nabla^{F_i}_V W - \dfrac{g(V,W)}{b_i} \operatorname{grad}_B b_i & \text{if } i = j, \end{cases}$

where $\nabla = \nabla^M$.

One can compute the *gradient* and the *Laplace-Beltrami* operator on M in terms of the *gradient* and the *Laplace-Beltrami* operator on B and F_i, respectively.

Proposition 2. *Let $M = B \times_{b_1} F_1 \times \cdots \times_{b_m} F_m$ be a pseudo-Riemannian multiply warped product with metric $g = g_B \oplus b_1^2 g_{F_1} \oplus \cdots \oplus b_m^2 g_{F_m}$ and $\phi \colon B \to \mathbb{R}$ and $\psi_i \colon F_i \to \mathbb{R}$ be smooth functions for any $i \in \{1, \cdots, m\}$. Then*

(1) $\operatorname{grad}(\phi \circ \pi) = \operatorname{grad}_B \phi$

(2) $\operatorname{grad}(\psi_i \circ \sigma_i) = \dfrac{1}{b_i^2} \operatorname{grad}_{F_i} \psi_i$

(3) $\Delta(\phi \circ \pi) = \Delta_B \phi + \displaystyle\sum_{i=1}^{m} s_i \frac{g_B(\operatorname{grad}_B \phi, \operatorname{grad}_B b_i)}{b_i}$

(4) $\Delta(\psi_i \circ \sigma_i) = \dfrac{\Delta_{F_i} \psi_i}{b_i^2}$

where $\Delta = \Delta_M$ *and* $\operatorname{grad} = \operatorname{grad}_M$.

Proposition 3. *Let* $M = B \times_{b_1} F_1 \times \cdots \times_{b_m} F_m$ *be a pseudo-Riemannian multiply warped product with metric* $g = g_B \oplus b_1^2 g_{F_1} \oplus \cdots \oplus b_m^2 g_{F_m}$ *also let* $X, Y, Z \in \mathfrak{L}(B)$ *and* $V \in \mathfrak{L}(F_i), W \in \mathfrak{L}(F_j)$ *and* $U \in \mathfrak{L}(F_k)$. *Then*

(1) $R(X,Y)Z = R_B(X,Y)Z$

(2) $R(V,X)Y = -\dfrac{\mathrm{H}_B^{b_i}(X,Y)}{b_i} V$

(3) $R(X,V)W = R(V,W)X = R(V,X)W = 0$ *if* $i \neq j$.

(4) $R(X,Y)V = 0$

(5) $R(V,W)X = 0$ *if* $i = j$.

(6) $R(V,W)U = 0$ *if* $i = j$ *and* $i, j \neq k$.

(7) $R(U,V)W = -g(V,W) \dfrac{g_B(\operatorname{grad}_B b_i, \operatorname{grad}_B b_k)}{b_i b_k} U$ *if* $i = j$ *and* $i, j \neq k$.

(8) $R(X,V)W = -\dfrac{g(V,W)}{b_i} \nabla_X^B (\operatorname{grad}_B b_i)$ *if* $i = j$.

(9) $R(V,W)U = R_{F_i}(V,W)U +$
$\dfrac{\| \operatorname{grad}_B b_i \|_B^2}{b_i^2} (g(V,U)W - g(W,U)V)$ *if* $i, j = k$.

where $R = R_M$.

Proposition 4. *Let* $M = B \times_{b_1} F_1 \times \cdots \times_{b_m} F_m$ *be a pseudo-Riemannian multiply warped product with metric* $g = g_B \oplus b_1^2 g_{F_1} \oplus \cdots \oplus b_m^2 g_{F_m}$, *also let* $X, Y, Z \in \mathfrak{L}(B)$ *and* $V \in \mathfrak{L}(F_i)$ *and* $W \in \mathfrak{L}(F_j)$. *Then*

(1) $\operatorname{Ric}(X,Y) = \operatorname{Ric}_B(X,Y) - \displaystyle\sum_{i=1}^{m} \frac{s_i}{b_i} \mathrm{H}_B^{b_i}(X,Y)$

(2) $\operatorname{Ric}(X,V) = 0$

(3) $\operatorname{Ric}(V,W) = 0$ *if* $i \neq j$.

(4) $\operatorname{Ric}(V,W) = \operatorname{Ric}_{F_i}(V,W) - \Big(\dfrac{\Delta_B b_i}{b_i} + (s_i - 1) \dfrac{\| \operatorname{grad}_B b_i \|_B^2}{b_i^2}$
$+ \displaystyle\sum_{k=1, k\neq i}^{m} s_k \frac{g_B(\operatorname{grad}_B b_i, \operatorname{grad}_B b_k)}{b_i b_k} \Big) g(V,W)$ *if* $i = j$,

where $\operatorname{Ric} = \operatorname{Ric}_M$.

3. Null Sectional Curvature

Suppose that Π is a null plane, that is Π consists of a one-dimensional subspace of null vectors and of space-like vectors perpendicular to that subspace. Let L be one of the null vectors and S be one of the space-like vectors. Since $Q(\Pi) = 0$, sectional curvature is not defined for a null plane. Then S. Harris introduces the *null sectional curvature* in [14] for degenerate planes as follows (see also [1]):

In order to define *null sectional curvature*, first we need to fix a choice of null vector.

Definition 4. *Let M be an n-dimensional Lorentzian manifold. A null congruence on M is a submanifold C of the tangent bundle T_0M of nonzero null vectors on M such that for all N in T_0M, there is exactly one scalar α satisfying $\alpha N \in C$.*

Then, *null sectional curvature* of Π with respect to N is given by

$$\mathrm{K}_N(\Pi) = \frac{g(R(L,S)S,L)}{g(S,S)}$$

where R is the Riemannian curvature tensor.

Remark 1. *Note that, this formula is independent of the choice of the space-like vector S in Π. However, $K_N(p,\Pi)$ depends quadratically on the null vector L.*

Null sectional curvature can be normalized via the help of a time-like vector field U in the following way: Assume that U is a time-like vector field on a Lorentzian manifold M. Then the null congruence $C(U)$ associated with U is given by

$$C(U) = \{L \in \mathrm{T_0M} | \mathrm{g(L,L)} = 0, \mathrm{g(L,U)} = -1\}.$$

Remark 2. *The null congruence lies in the future null cone due to the -1 in the definition.*

Then U-normalized null sectional curvature is

$$\mathrm{K}_U(\Pi) = \frac{g(R(L,S)S,L)}{g(S,S)}$$

where

$$C_U(M) = \{L \in TM |\, g(L,L) = 0 \quad \text{and} \quad g(L, U_{\Pi(L)}) = -1\}.$$

3.1. **Null Sectional Curvature of a multiply GRW space-time.** Let $M = I \times_{b_1} F_1 \ldots \times_{b_m} F_m$ be a multiply GRW space-time with the metric $g = -\mathrm{d}t^2 \oplus b_1^2 g_{F_1} \oplus \ldots \oplus b_m^2 g_{F_m}$. Let $L = -\partial_t + \mathbf{V}$ and $S = Y + \mathbf{W}$ where $\mathbf{V} = \mathbf{\Sigma} V_i$ $\mathbf{W} = \mathbf{\Sigma} W_j$. Then,

$$
\begin{aligned}
g(R(L,S)S,L) &= g(R(-\partial_t+\mathbf{V},Y+\mathbf{W})\mathbf{V},Y+\mathbf{W},-\partial_t+\mathbf{V})\\
&= g(R(-\partial_t,Y)Y,-\partial_t)+g(R(\mathbf{\Sigma}V_i,Y)Y,-\partial_t)\\
&+ g(R(-\partial_t,\mathbf{\Sigma}W_j)Y,-\partial_t)+g(R(\mathbf{\Sigma}V_i,\mathbf{\Sigma}W_j)Y,-\partial_t)\\
&+ g(R(-\partial_t,Y)\mathbf{\Sigma}W_j,-\partial_t)+g(R(\mathbf{\Sigma}V_i,Y)\mathbf{\Sigma}W_j,-\partial_t)\\
&+ g(R(-\partial_t,\mathbf{\Sigma}W_j)\mathbf{\Sigma}W_j,-\partial_t)+g(R(\mathbf{\Sigma}V_i,\mathbf{\Sigma}W_j)\mathbf{\Sigma}W_j,-\partial_t)\\
&+ g(R(-\partial_t,Y)Y,\mathbf{\Sigma}V_i)+g(R(\mathbf{\Sigma}V_i,Y)Y,\mathbf{\Sigma}V_i)\\
&+ g(R(-\partial_t,\mathbf{\Sigma}W_j)Y,\mathbf{\Sigma}V_i)+g(R(\mathbf{\Sigma}V_i,\mathbf{\Sigma}W_j)Y,\mathbf{\Sigma}V_i)\\
&+ g(R(-\partial_t,Y)\mathbf{\Sigma}W_j,\mathbf{\Sigma}V_i)+g(R(\mathbf{\Sigma}V_i,Y)\mathbf{\Sigma}W_j,\mathbf{\Sigma}V_i)\\
&+ g(R(-\partial_t,\mathbf{\Sigma}W_j)\mathbf{\Sigma}W_j,\mathbf{\Sigma}V_i)+g(R(\mathbf{\Sigma}V_i,\mathbf{\Sigma}W_j)\mathbf{\Sigma}W_j,\mathbf{\Sigma}V_i)
\end{aligned}
$$

$$
\begin{aligned}
g(R(\mathbf{V},\mathbf{W})\mathbf{W}),\mathbf{V}) &= \mathbf{\Sigma}_{i\neq j}-\frac{1}{b_j^2}b_j''g(W_j,W_j)g(V_i,V_i)\\
&+ \mathbf{\Sigma}g(R_{F_i}(V_i,W_i),W_i,V_i)\\
&+ \mathbf{\Sigma}\frac{1}{b_i^2}b_i''g(g(V_i,W_i)W_i-g(W_i,W_i)V_i),V_i)\\
&= \mathbf{\Sigma}_{i\neq j}-(b_j')^2g_{F_j}(W_j,W_j)b_i^2g_{F_i}(V_i,V_i)\\
&+ \mathbf{\Sigma}b_i^2g_{F_i}(R_{F_i}(V_i,W_i),W_i,V_i)\\
&+ \mathbf{\Sigma}b_i^2(b_i')^2[g_{F_i}(V_i,W_i)^2-g_{F_i}(V_i,V_i)g_{F_i}(W_i,W_i)]
\end{aligned}
\tag{3.1}
$$

Theorem 1. *Let $M=I\times_{b_1}F_1\ldots\times_{b_m}F_m$ be a multiply GRW space-time with the metric $g=-\mathrm{d}t^2\oplus b_1^2g_{F_1}\oplus\ldots\oplus b_m^2g_{F_m}$. Assume that $L=-\partial_t+\mathbf{V}$ and $S=Y+\mathbf{W}$ where $\mathbf{V}=\mathbf{\Sigma}V_i$ and $\mathbf{W}=\mathbf{\Sigma}W_j$. Then null sectional of (M,g) is given by*

$$
\begin{aligned}
\mathrm{P}(p,\Pi) &= \mathbf{\Sigma}b_kg_{F_k}(W_k,V_k)H^B_{b_k}(\partial_t,Y)\\
&+ \mathbf{\Sigma}b_kg_{F_k}(V_k,V_k)\mathrm{H}^B_{b_k}(Y,Y)\\
&- \mathbf{\Sigma}b_jb_j''g_{F_j}(W_j,W_j)\\
&+ \mathbf{\Sigma}b_ig_{F_i}(W_i,V_i)\mathrm{H}^{b_i}_B(\partial_t,Y)\\
&- \mathbf{\Sigma}_{j\neq k}b_k(b_j')^2g_{F_k}(V_k,V_k)g_{F_j}(W_j,W_j)\\
&+ \mathbf{\Sigma}b_i^2g_{F_i}(R_{F_i}(W_i,V_i)V_i,W_i)\\
&- \mathbf{\Sigma}b_i^2(b_i')^2b_i''[(g_{F_i}(W_i,V_i))^2-g_{F_i}(V_i,V_i)g_{F_i}(W_i,W_i)]
\end{aligned}
$$

$$
\begin{aligned}
g(S,S) &= g(Y+\mathbf{\Sigma}W_j,Y+\mathbf{\Sigma}W_j)\\
&= g_B(Y,Y)+\mathbf{\Sigma}(b_j)^2g_{F_j}(W_j,W_j)
\end{aligned}
$$

By taking Y as $h\partial_t$ in Theorem 1, we obtain the following result.

Corollary 1. *Let $M = I \times_{b_1} F_1 \ldots \times_{b_m} F_m$ be a MGRW space-time with the metric $\mathrm{g} = -\mathrm{d}t^2 \oplus b_1^2 g_{F_1} \oplus \ldots \oplus b_m^2 \mathrm{g}_{F_m}$. Suppose that $L = -\partial_t + \mathbf{V}$ and $S = h\partial_t + \mathbf{W}$ where $\mathbf{V} = \Sigma V_i$ and $\mathbf{W} = \Sigma W_j$ and $h : I \to \mathbb{R}$ is a smooth function. Then,*

$$\begin{aligned}
\mathrm{P}(p,\Pi) &= \mathbf{\Sigma} h b_k b_k'' \mathrm{g}_{F_k}(W_k, V_k) \\
&+ \mathbf{\Sigma} h^2 b_k b_k'' \mathrm{g}_{F_k}(V_k, V_k) \\
&- \mathbf{\Sigma} b_j b_j'' \mathrm{g}_{F_j}(W_j, W_j) \\
&+ \mathbf{\Sigma} h b_i b_i'' \mathrm{g}_{F_i}(W_i, V_i) \\
&- \mathbf{\Sigma}_{j\neq k} b_k (b_j')^2 \mathrm{g}_{F_k}(V_k, V_k) \mathrm{g}_{F_j}(W_j, W_j) \\
&+ \mathbf{\Sigma} b_i^2 \mathrm{g}_{F_i}(\mathrm{R}_{F_i}(W_i, V_i)V_i, W_i) \\
&- \mathbf{\Sigma} b_i (b_i')^4 b_i'' [(\mathrm{g}_{F_i}(W_i, V_i))^2 - \mathrm{g}_{F_i}(V_i, V_i)\mathrm{g}_{F_i}(W_i, W_i)]
\end{aligned}$$

$$\begin{aligned}
g(S,S) &= g(h\partial t + \mathbf{\Sigma} W_j, h\partial t + \mathbf{\Sigma} W_j) \\
&= -h'' + \mathbf{\Sigma} (b_j)^2 g_{F_j}(W_j, W_j)
\end{aligned}$$

3.2. **Null Sectional Curvature of a GRW space-time.** Assume that $M = I \times_b F$ is a GRW space-time with the metric $g = g_I \oplus b^2 g_F$ where $g_I = -dt^2$. Let Π be degenerate null plane at $p \in M$ spanned by a null vector L and space-like vector S,i.e., $g(L,L) = 0$ and $g(S,S) > 0$ where $U = \partial_t$ is a reference frame since $g(U,U) = g(\partial_t, \partial_t) = -1$.

(i) Let $L = h\partial_t + V$, $U = \partial_t$
$g(L,U) = g(h\partial_t + V, \partial_t) = -dt^2(h\partial_t, \partial_t) + b^2 g_F(V, 0)$
$= -h$
This implies $g(L,U) = 1$ if and only if $h = -1$. Then we have $L = -\partial_t + V$

(ii) $g(L,L) = g(-\partial_t + V, -\partial_t + V) = -dt^2(-\partial_t, V) + b^2 g_F(V,V) = -1 + b^2 g_F(V,V) = -1 + g_F(V,V)$
By imposing L to be null, i.e, $g(L,L) = 0$, we obtain $g_F(V,V) = \frac{1}{b^2}$.

(iii) Let $E = \mathrm{span}(\{L, S\})$ be a degenerate plane section, i.e., $Q(L,S) = 0$
$Q(L,S) = Q(-\partial_t + V, Y + W) = g(-\partial_t, -\partial_t)g(Y+W, Y+W) - g(-\partial_t, Y+W)^2 = [-g_I(\partial_t, Y) + b^2 g_F(V,W)]^2 = 0$ This implies $g_F(V,W) = \frac{1}{b^2} g_I(\partial_t, Y)$

Corollary 2. *Let $M = I \times_b F$ be a GRW space-time with the metric $g = -dt^2 \oplus b^2 g_F$ where $b : I \to (0,\infty)$ is a smooth function. Then, null sectional curvature of (M,g) is as follows:*

$$\begin{aligned}
g(R(L,S)S,L) &= -bb'' g_F(W,W) \\
&+ b^2 g_F(R_F(W,V)V,W)) \\
&+ b g_F(V,V) \mathrm{H}_B^b(Y,Y) \\
&+ b^2 (b')^2 [g_F(V,W)^2 - \frac{1}{b^2} g_F(W,W)]
\end{aligned} \tag{3.2}$$

$$g(S,S) = g_I(Y,Y) + b^2 g_F(W,W) \tag{3.3}$$

Remark 3. *As a special case assume that $Y = 0$. Then*

$$\begin{aligned} \mathrm{Q}_F(V,W) &= g_F(V,V)g_F(W,W) - g_F(V,W)^2 \\ &= \frac{1}{b^2} g_F(W,W) \end{aligned} \tag{3.4}$$

In this case we have,

$$\mathrm{K}_U(\Pi) = \frac{1}{b^2} K_F(V,W) + \frac{b''}{b} - (\frac{b'}{b})^2 \tag{3.5}$$

Using the above equation we establish that $\mathrm{K}_U(\Pi) = \frac{1}{b^2}\mathrm{K}_F(V,W)$ if and only if the warping function $b(t) = ce^{kt}$ where c and k are arbitrary constants.

3.3. **Null Sectional Curvature of a Generalized Kasner Space-time.**

Definition 5. *A generalized Kasner space-time (M,g) is a Lorentzian multiply warped product of the form $M = I \times_{\varphi^{p_1}} F_1 \ldots \times_{\varphi^{p_m}} F_m$ with the metric $\mathrm{g} = -\mathrm{d}t^2 \oplus \varphi^{2p_1} g_{F_1} \times \oplus \ldots \oplus \varphi^{2p_m} \mathrm{g}_{F_m}$ where $\varphi : I \to (0,\infty)$ is smooth and $p_i \in \mathbb{R}$ for $i = 1,\ldots,m$ and also $I = (t_1,t_2)$ with $-\infty \le t_1 < t_2 \le \infty$.*

Corollary 3. *Let $L = -\partial_t + \mathbf{V}$ and $S = Y + \mathbf{W}$ where $\varphi \to (0,\infty)$ with $-\infty \le t_1 < t_2 \le \infty$. and $\mathbf{V} = \mathbf{\Sigma} V_i$, $\mathbf{W} = \mathbf{\Sigma} W_j$ Then null sectional of a multiply Kasner space-time of the form above is given by*

$$\begin{aligned} \mathrm{P}(p,\Pi) = {} & \mathbf{\Sigma}\varphi^{p_k}\mathrm{g}_{F_k}(W_k,V_k)\mathrm{H}_I^{\varphi^{p_k}}(\partial_t,Y) \\ & + \mathbf{\Sigma}\varphi^{p_k}\mathrm{g}_{F_k}(V_k,V_k)\mathrm{H}_I^{\varphi^{p_k}}(Y,Y) \\ & - \mathbf{\Sigma}\varphi^{p_j} p_j(p_j-1)\varphi^{(p_j-2)}\mathrm{g}_{F_j}(W_j,W_j) \\ & + \mathbf{\Sigma}\varphi^{p_i}\mathrm{g}_{F_i}(W_i,V_i)\mathrm{H}_I^{\varphi^{p_i}}(\partial_t,Y) \\ & - \mathbf{\Sigma}_{j\neq k}\varphi^{p_k} {p_j}^2 \varphi^{2(p_j-1)}\mathrm{g}_{F_k}(V_k,V_k)\mathrm{g}_{F_j}(W_j,W_j) \\ & + \mathbf{\Sigma}\varphi^{2p_i}\mathrm{g}_{F_i}(\mathrm{R}_{F_i}(W_i,V_i)V_i,W_i) \\ & - \mathbf{\Sigma}\varphi^{2p_i} p_i^2 \varphi^{2(p_i-1)} p_i(p_i-1)\varphi^{(p_i-2)}[(\mathrm{g}_{F_i}(W_i,V_i))^2 - \mathrm{g}_{F_i}(V_i,V_i)\mathrm{g}_{F_i}(W_i,W_i)] \end{aligned}$$

$$\begin{aligned} \mathrm{g}(S,S) &= \mathrm{g}(Y + \mathbf{\Sigma}W_j, Y + \mathbf{\Sigma}W_j) \\ &= \mathbf{\Sigma}\varphi^{2p_j}\mathrm{g}_I(Y,Y) + \mathrm{g}_{F_j}(W_j,W_j) \end{aligned}$$

4. Four-dimensional Space-time Models

4.1. **Type I: Null Sectional Curvature of a Kasner Space-time with fiber of dimension** (3)**.**

Corollary 4. *Let $M = I \times_b F$ and $g = -dt^2 \oplus b^2 g_F$, $L = -\partial_t + V$ and $S = h\partial_t + W$ where Π be a degenerate null plane which is spanned by the tangent vectors L and S. Then,*

$$\begin{aligned} P(\Pi) = & f^2 bb'' g_F(V,V) \\ & - bb'' g_F(W,W) \\ & + bb'' g_F(V,W) \\ & + b^2 g_F(R_F(W,V)V,W) \\ & - b^2 (b')^2 [g_F(V,W)^2 - g_F(V,V) g_F(V,W)] \end{aligned} \tag{4.1}$$

4.2. **Type II: Null Sectional Curvature of a Kasner Space-time with fiber of dimension** $(1,2)$**.** A Kasner space-time with fiber of dimension $(1,2)$ is a Lorentzian multiply warped product (M,g) of the form $M = (0,\infty) \times_{b_1} \mathbb{R} \times_{b_2} F$ with the metric $g = -dt^2 \oplus b_1^2 dx^2 \oplus b_2^2 g_F$

Corollary 5. *Let Π be degenerate null plane at $p \in M$ spanned by a null vector L and space-like vector S, i.e., $g(L,L) = 0$ and $g(S,S) > 0$.*

Let $L = -\partial t + f_1 \partial x + V$ and $S = f\partial t + h_1 \partial x + W$ where

$$V_1 = f_1 \partial_x,$$

$$W_1 = h_1 \partial_x$$

Then the null sectional curvature of Π with respect to L is is given by

$$K_U(\Pi) = \frac{R((L,S)S,L)}{g(S,S)}$$

where $U = \partial_t$ is a reference frame since $g(U,U) = g(\partial_t, \partial_t) = -1$ and

$$\begin{aligned} g(R(\Sigma V_i, \Sigma W_j)\Sigma W_j, \Sigma V_i) = & b_1 f_1 h_1 f b_1'' + b_2 f^2 b_2'' g_F(V,V) \\ & - b_1 b_1'' h_1^2 - b_2 b_2'' g_F(W,W) \\ & + b_1 f_1 h_1 f b_1'' + b_2 b_2'' g_F(V,W) \\ & + b_2^2 g_F(R_F(W,V)V,W) \\ & - b_2^2 (b_2')^2 [g_F(V,W)^2 \\ & - g_F(V,V) g_F(W,W)] \end{aligned} \tag{4.2}$$

$$\begin{aligned} g(S,S) &= g(f\partial t + \Sigma W_j, f\partial t + \Sigma W_j) \\ &= -f^2 + b_1^2 h_1^2 + b_2^2 g_F(W,W) \end{aligned} \tag{4.3}$$

Remark 4. *For the case $Y = h\partial_t$ and $V_1 = W_1 = 0$, $P(E)$ becomes as follows:*

$$\begin{aligned} P(\Pi) &= \varphi^{p_2} p_2(p_2-1)\varphi^{p_2-2} g_{F_2}(W_2, V_2) \\ &+ h^2 \varphi^{p_2} p_2(p_2-1)\varphi^{p_2-2} g_{F_2}(V_2, V_2) \\ &- \varphi^{p_2} p_2(p_2-1)\varphi^{p_2-2} g_{F_2}(W_2, V_2) \\ &+ h\varphi^{p_2} p_2(p_2-1)\varphi^{p_2-2} g_{F_2}(W_2, V_2) \\ &+ \varphi^{2p_2} g_{F_2}(R_{F_2}(W_2, V_2)V_2, W_2) \\ &- p_2^4 \varphi^{6p_2-4}[g_{F_2}(W_2, V_2)^2 - g_{F_2}(V_2, V_2) g_{F_2}(W_2, W_2)] \end{aligned} \tag{4.4}$$

Similarly we can obtain $P(E)$ for the case $V_2 = W_1 = 0$ by using the symmetry properties.

4.3. **Type III: Null Sectional Curvature of a Kasner Space-time with the fiber of dimension** $(1, 1, 1)$**.** A Kasner space-time with the fiber of dimension $(1, 1, 1)$ is a Lorentzian multiply warped product (M, g) of the form

$$M = (0, \infty) \times_{\varphi^{p_1}} \mathbb{R} \times_{\varphi^{p_2}} \mathbb{R} \times_{\varphi^{p_3}} \mathbb{R}$$

with the metric

$$g = -dt^2 \oplus \varphi^{2p_1} dx^2 \oplus \varphi^{2p_2} dy^2 \oplus \varphi^{2p_3} dz^2$$

where $p_1 + p_2 + p_3 = p_1^2 + p_2^2 + p_3^2 = 1$.

Corollary 6. *Let Π be degenerate null plane at $p \in M$ spanned by a null vector L and spacelike vector S,i.e., $g(L, L) = 0$ and $g(S, S) > 0$.*

Let $L = -\partial_t + \Sigma V_i$ and $S = f\partial_t + \Sigma W_j$ where

$$V_1 = f_1\partial_x, V_2 = f_2\partial_y$$

$$V_3 = f_3\partial_z, W_1 = h_1\partial_x$$

$$W_2 = h_2\partial_y, W_3 = h_3\partial_z$$

Then the null sectional curvature of Π with respect to L is is given by

$$K_U(\Pi) = \frac{R((L, S)S, L)}{g(S, S)}$$

where $U = \partial_t$ is a reference frame since $g(U, U) = g(\partial_t, \partial_t) = -1$.

$$\begin{aligned} g(R(\Sigma V_i, \Sigma W_j)\Sigma W_j, \Sigma V_i) &= -\Sigma \varphi^{p_i} f_i h_i \\ &+ \Sigma \varphi^{p_k} (f_k)^2 f^2 p_k(p_k-1)\varphi^{p_k-2} \\ &- \Sigma p_j(p_j-1)\varphi^{p_j-2}(\varphi^{p_j} h_j^2 \\ &+ \Sigma \varphi^{p_i} f_i h_i f p_i(p_i-1)\varphi^{p_i-2} \\ &- \Sigma_{j\neq k} \varphi^{p_k} f_k^2 p_j^2 \varphi^{(2p_j-2)} h_j^2 \end{aligned} \tag{4.5}$$

$$
\begin{aligned}
g(S,S) &= g(f\partial t + \Sigma W_j, f\partial t + \Sigma W_j) \\
&= -f^2 \Sigma \varphi^{2p_j} h_j^2
\end{aligned} \tag{4.6}
$$

5. Null sectional curvature of a SSS-T

Let $M = f_I \times F$ be a SSST with the metric $\mathrm{g} = -f^2 \mathrm{g}_I \oplus \mathrm{g}_F$ where $-\mathrm{d}t^2$ is the negative definite metric on I.

Let Π be degenerate null plane at $p \in M$ spanned by a null vector L and spacelike vector S,i.e., $\mathrm{g}(L,L) = 0$ and $\mathrm{g}(S,S) > 0$ with the reference frame $U = f^{-1}\partial_t$. The following calculations will be useful in Corollary 7:

(i) Let $L = h\partial_t + V$, $U = f^{-1}\partial_t$
$g(L,U) = \mathrm{g}(h\partial_t + V, f^{-1}\partial_t) = -f^2 dt^2(h\partial_t, f^{-1}\partial_t) + \mathrm{g}_F(V,0)$
$= -f^2 h f^{-1}$
$= -fh$
This implies $\mathrm{g}(L,U) = 1$ iff $h = -f^{-1}$. Then we have $L = -f^{-1}\partial_t + V$

(ii) $\mathrm{g}(L,L) = \mathrm{g}(-f^{-1}\partial_t + V, -f^{-1}\partial_t + V) = -f^2 dt^2(-f^{-1}\partial_t, v) + \mathrm{g}_F(V,V) = -f^2 \frac{1}{f^2} + \mathrm{g}_F(V,V) = -1 + \mathrm{g}_F(V,V)$
If we want to make $g(L,L) = 0$, i.e., L is null, then we must have $g_F(V,V) = 1$.

(iii) Let $E = \mathrm{span}(\{L,S\})$ be a degenerate plane section,i.e., $Q(L,S) = 0$
$Q(L,S) = Q(-f^{-1}\partial_t + V, Y + W) = \mathrm{g}(-f^{-1}\partial_t, -f^{-1}\partial_t)\mathrm{g}(Y+W, Y+W) - \mathrm{g}(-f^{-1}\partial_t, Y+W)^2 = [-f\mathrm{g}_I(\partial_t, Y) + \mathrm{g}_F(V,W)]^2 = 0$ This implies $\mathrm{g}_F(V,W) = f\mathrm{g}_I(\partial_t, Y)$

Corollary 7. *Then the null sectional curvature of SSST is as follows:*

$$
\begin{aligned}
\mathrm{g}(\mathrm{R}(L,S)S,L) = &-\mathrm{g}_F(\nabla_F f, \nabla_F f)(\mathrm{g}_I(Y,\partial_t)^2 + \mathrm{g}_I(Y,Y)) \\
&- f\mathrm{g}_I(Y,Y)\mathrm{H}_F^f(V,V) \\
&- \mathrm{g}_I(Y,\partial_t)\mathrm{H}_F^f(V,W) \\
&+ \mathrm{g}_I(Y,\partial_t)\mathrm{H}_F^f(V,W) \\
&+ \frac{1}{f}\mathrm{H}_F^f(W,W) \\
&+ \mathrm{g}_F(R_F(V,W)V,W)
\end{aligned} \tag{5.1}
$$

If we take $Y = h\partial_t$, we obtain that:

$$
\begin{aligned}
\mathrm{g}(\mathrm{R}(L,S)S,L) = &\, fh^2\mathrm{H}_F^f(V,V) \\
&+ \frac{1}{f}\mathrm{H}_F^f(W,W) + \mathrm{g}_F(R_F(V,W)V,W)
\end{aligned} \tag{5.2}
$$

Since null sectional curvature is independent from the choice of S (space-like vector), without loss of generality we can assume that $\mathrm{g}(S,S)=1$. Then

$$\begin{aligned}\mathrm{g}(\mathrm{R}(L,S)S,L) &= -\mathrm{g}_F(\nabla_F f,\nabla_F f)(\mathrm{g}_I(Y,\partial_t)^2+\mathrm{g}_I(Y,Y))\\ &\quad - f\mathrm{g}_I(Y,Y)\mathrm{H}^f_F(V,V)\\ &\quad - \frac{1}{f}\mathrm{H}^f_F(W,W)\\ &\quad + \mathrm{g}_F(R_F(V,W)V,W)\end{aligned} \tag{5.3}$$

Remark 5. *As a special case assume that $Y=0$. Then*

$$Q_F(V,W)=\mathrm{g}_F(W,W) \tag{5.4}$$

In this case we have

$$K_U(\Pi)=K_F(V,W)-\frac{H^f_F(W,W)}{f\mathrm{g}_F(W,W)} \tag{5.5}$$

If we assume that $\mathrm{H}^f_F=kf\mathrm{g}_F(W,W)$ for some $k\in\mathbb{R}$, then

$$K_U(\Pi)=K_F(V,W)-k$$

Appendix A. Null Sectional Curvature of MGRW space-time

Here we give main steps of the proof of Theorem 1:

$$\begin{aligned}g(R(L,S)S,L) &= g(R(-\partial_t+\mathbf{\Sigma}V_i,Y+\mathbf{\Sigma}W_j)Y+\mathbf{\Sigma}W_j,-\partial_t+\mathbf{\Sigma}V_i)\\ &= g(R(-\partial_t,Y)Y,-\partial_t)+g(R(\mathbf{\Sigma}V_i,Y)Y,-\partial_t)\\ &\quad + g(R(-\partial_t,\mathbf{\Sigma}W_j)Y,-\partial_t)+g(R(\mathbf{\Sigma}V_i,\mathbf{\Sigma}W_j)Y,-\partial_t)\\ &\quad + g(R(-\partial_t,Y)\mathbf{\Sigma}W_j,-\partial_t)+g(R(\mathbf{\Sigma}V_i,Y)\mathbf{\Sigma}W_j,-\partial_t)\\ &\quad + g(R(-\partial_t,\mathbf{\Sigma}W_j)\mathbf{\Sigma}W_j,-\partial_t)+g(R(\mathbf{\Sigma}V_i,\mathbf{\Sigma}W_j)\mathbf{\Sigma}W_j,-\partial_t)\\ &\quad + g(R(-\partial_t,Y)Y,\mathbf{\Sigma}V_i)+g(R(\mathbf{\Sigma}V_i,Y)Y,\mathbf{\Sigma}V_i)\\ &\quad + g(R(-\partial_t,\mathbf{\Sigma}W_j)Y,\mathbf{\Sigma}V_i)+g(R(\mathbf{\Sigma}V_i,\mathbf{\Sigma}W_j)Y,\mathbf{\Sigma}V_i)\\ &\quad + g(R(-\partial_t,Y)\mathbf{\Sigma}W_j,\mathbf{\Sigma}V_i)+g(R(\mathbf{\Sigma}V_i,Y)\mathbf{\Sigma}W_j,\mathbf{\Sigma}V_i)\\ &\quad + g(R(-\partial_t,\mathbf{\Sigma}W_j)\mathbf{\Sigma}W_j,\mathbf{\Sigma}V_i)+g(R(\mathbf{\Sigma}V_i,\mathbf{\Sigma}W_j)\mathbf{\Sigma}W_j,\mathbf{\Sigma}V_i)\end{aligned}$$

$$\begin{aligned}g(R(-\partial_t,Y)Y,-\partial_t) &= g_I(R_I(\partial_t,Y)Y,-\partial_t)\\ &= 0\end{aligned} \tag{A.1}$$

$$\begin{aligned} g(R(\mathbf{\Sigma}V_i, Y)Y, -\partial_t) &= \mathbf{\Sigma} g(-\frac{1}{b_i}\mathrm{H}_I^{b_i}(Y,Y)V_i, -\partial_t) \\ &= \mathbf{\Sigma}\frac{1}{b_i}\mathrm{H}_B^{b_i}(Y,\partial_t)g(V_i, -\partial_t) \\ &= 0 \end{aligned} \tag{A.2}$$

$$\begin{aligned} g(R(-\partial_t, \mathbf{\Sigma}W_j)Y, -\partial_t) &= \mathbf{\Sigma} g(-\frac{1}{b_j}\mathrm{H}_I^{b_j}(-\partial_t, Y)W_j, -\partial_t) \\ &= \Sigma\frac{1}{b_j}\mathrm{H}_I^{b_j}(\partial_t, Y)g(W_j, \partial_t) \\ &= 0 \end{aligned} \tag{A.3}$$

$$g(R(\mathbf{\Sigma}V_i, \mathbf{\Sigma}W_j)Y, -\partial_t) = 0 \tag{A.4}$$

$$g(R(-\partial_t, Y)\mathbf{\Sigma}W_j, -\partial_t) = 0 \tag{A.5}$$

$$\begin{aligned} g(R(\mathbf{\Sigma}V_i, Y)\mathbf{\Sigma}W_j, -\partial_t) &= \mathbf{\Sigma} g(\frac{-1}{b_i}g(V_i, W_i)\nabla_Y^I(\nabla^I b_i), -\partial_t) \\ &= \mathbf{\Sigma}\frac{-b_i^2}{b_i}g_{F_i}(V_i, W_i)g(\nabla_Y^I(\nabla^I b_i), -\partial_t) \\ &= \mathbf{\Sigma} b_i g_{F_i}(V_i, W_i)\mathrm{H}_I^{b_i}(Y, \partial_t) \end{aligned} \tag{A.6}$$

$$\begin{aligned} g(R(-\partial_t, \mathbf{\Sigma}W_j)\mathbf{\Sigma}W_j, -\partial_t) &= \mathbf{\Sigma} g(\frac{-1}{b_j}g(V_i, W_i)\nabla_{-\partial_t}^I(\nabla^I b_i), Y) \\ &= \mathbf{\Sigma}\frac{-1}{b_i}g_{F_j}(W_j, W_j)g(\nabla_{-\partial_t}^I(\nabla^I b_j), -\partial_t) \\ &= \mathbf{\Sigma} - b_j g_{F_j}(W_j, W_j)\mathrm{H}_I^{b_i}(-\partial_t, -\partial_t) \\ &= \mathbf{\Sigma} - b_j b_j'' g_{F_j}(W_j, W_j) \end{aligned} \tag{A.7}$$

(A.8)

$$\begin{aligned} g(R(\mathbf{\Sigma}V_i, \mathbf{\Sigma}W_j)\mathbf{\Sigma}W_j, -\partial_t) &= \mathbf{\Sigma}_{i\neq j} g(g(W_j, W_j)\frac{g_I(\nabla^I b_j, \nabla^I b_i)}{b_i b_j}V_i, \partial_t) \\ &+ g(R_{F_i}(V_i, W_i)W_i, -\partial_t) \\ &+ \frac{g_I(\nabla^I b_i, \nabla^I b_i)}{b_i b_i}g(g(V_i, W_i)W_i - g(W_i, W_i)V_i, -\partial_t) \\ &= 0 \end{aligned}$$

$$\begin{aligned} g(R(-\partial_t, Y)Y, \mathbf{\Sigma}V_i) &= g(R_I(\partial_t, Y)Y, \mathbf{\Sigma}V_i) \\ &= 0 \end{aligned} \tag{A.9}$$

$$
\begin{aligned}
g(R(\mathbf{\Sigma}V_i, Y)Y, \mathbf{\Sigma}V_i) &= \mathbf{\Sigma}g(-\frac{\mathrm{H}_I^{b_i}(Y, -\partial_t)}{b_i}V_i, V_i) \\
&= \mathbf{\Sigma}b_i^2\frac{\mathrm{H}_B^{b_i}(Y, Y)}{b_i}g_{F_i}(V_i, V_i) \\
&= \mathbf{\Sigma}b_i\mathrm{H}_B^{b_i}(Y, Y)g_{F_i}(V_i, V_i)
\end{aligned} \tag{A.10}
$$

$$
\begin{aligned}
g(R(-\partial_t, \mathbf{\Sigma}W_j)Y, \mathbf{\Sigma}V_i) &= \mathbf{\Sigma}g(-\frac{\mathrm{H}_I^{b_i}(-\partial_t, Y)}{b_i}W_j, V_i) \\
&= \mathbf{\Sigma}b_i^2\frac{\mathrm{H}_I^{b_i}(\partial_t, Y)}{b_i}g_{F_i}(V_i, W_i) \\
&= \mathbf{\Sigma}b_i\mathrm{H}_I^{b_i}(\partial_t, Y)g_{F_i}(V_i, W_i)
\end{aligned} \tag{A.11}
$$

$$
g(R(\mathbf{\Sigma}V_i, \mathbf{\Sigma}W_j)Y, \mathbf{\Sigma}V_i) = 0 \tag{A.12}
$$

$$
g(R(-\partial_t, Y)\mathbf{\Sigma}W_j, \mathbf{\Sigma}V_i) = 0 \tag{A.13}
$$

$$
\begin{aligned}
g(R(\mathbf{\Sigma}V_i, Y)\mathbf{\Sigma}W_j, \mathbf{\Sigma}V_i) &= \mathbf{\Sigma}g(\frac{g(V_i, W_i)}{b_i}\nabla_Y^I(\nabla^I b_i), V_i) \\
&= \mathbf{\Sigma}b_i g_{F_i}(V_i, W_i)g(\nabla_Y^I(\nabla^I b_i), V_i) \\
&= \mathbf{\Sigma}\frac{-g_{F_i}(V_i, W_i)}{b_i}\mathrm{H}_{b_i}^I(Y, V_i) \\
&= 0
\end{aligned} \tag{A.14}
$$

$$
\begin{aligned}
g(R(-\partial_t, \mathbf{\Sigma}W_j)\mathbf{\Sigma}W_j, \mathbf{\Sigma}V_i) &= \mathbf{\Sigma}g(\frac{g(W_j, W_j)}{b_j}\nabla_{-\partial_t}^I(\nabla^I b_i), V_i) \\
&= \mathbf{\Sigma} - b_j g_{F_j}(W_j, W_j)g(\nabla_{\partial_t}^I(\nabla^I b_i), V_i)) \\
&= 0
\end{aligned} \tag{A.15}
$$

(A.16)

$$\begin{aligned} g(R(\mathbf{\Sigma}V_i, \mathbf{\Sigma}W_j)\mathbf{\Sigma}W_j, \mathbf{\Sigma}V_i) &= \mathbf{\Sigma}_{i\neq j} g(-g(W_j, W_j)\frac{g_I(\nabla^I b_j, \nabla^I b_j)}{b_j^2}V_i, V_i) \\ &+ \mathbf{\Sigma} g(R_{F_i}(V_i, W_i), W_i, V_i) \\ &+ g(\frac{g_I(\nabla^I b_i, \nabla^I b_i)}{b_i^2}(g(V_i, W_i)W_i - g(W_i, W_i)V_i), V_i) \\ &= \mathbf{\Sigma}_{i\neq j} - (b_j')^2 g_{F_j}(W_j, W_j) b_i^2 g_{F_i}(V_i, V_i) \\ &+ \mathbf{\Sigma} b_i^2 g_{F_i}(R_{F_i}(V_i, W_i), W_i, V_i) \\ &+ \mathbf{\Sigma} b_i^2 (b_i')^2 [g_{F_i}(V_i, W_i)^2 - g_{F_i}(V_i, V_i) g_{F_i}(W_i, W_i)] \end{aligned}$$

Then we obtain the following formula for null sectional curvature of Multiply Generalized Robertson Walker Spacetimes in Theorem 1.

$$\begin{aligned} g(S, S) &= g(Y + \mathbf{\Sigma}W_j, Y + \mathbf{\Sigma}W_j) \\ &= \mathbf{\Sigma}(b_j)^2 g_B(Y, Y) g_{F_j}(W_j, W_j) \end{aligned}$$

References

[1] A. L. Albujer and S. Haesen, *A geometrical interpretation of the null sectional curvature*, J. Geom. Phys., **60**(3), (2010), 471476.

[2] D. E. Allison, *Energy conditions in standard static space-times*, General Relativity and Gravitation, **20**(2), (1988), 115–122.

[3] D. E. Allison, *Geodesic completeness in static spacetimes*, Geometriae Dedicata **26**, (1988), 85-97.

[4] J. K. Beem, J. K. and P. E. Ehrlich, Th. G. Powell, *Warped product manifolds in relativity*, Selected Studies: A Volume Dedicated to the Memory of Albert Einstein, North-Holland, Amsterdarm (1982), 41–56

[5] J. K. Beem and P. E. Ehrlich and K. L. Easley, *Global Lorentzian Geometry*, (2nd Ed.), Marcel Dekker, New York, (1996).

[6] A. L. Besse, *Einstein Manifolds*, Springer-Verlag, Heidelberg, (2008).

[7] R. L. Bishop and B. O'Neill, *Manifolds of Negative Curvature*, Transections of the American Mathematical Society (1969), 1-49.

[8] M. Dajczer and K. Nomizu, *On the boundness of Ricci curvature of an indefinite metric*, Bol. Soc. Bras. Mat., **11**(1), (1980), 25-30.

[9] M. Dajczer, Marcos and K. Nomizu, *On Sectional Curvature of Indefinite Metrics, II*, Math. Ann. **247**(3), (1980), 279-282.

[10] F. Dobarro and B. Ünal, *Curvature of Multiply Warped Products*, J. Geom. Phys., **55**(1), (2005), 75-106.

[11] L. Graves and K. Nomizu, *On sectional curvature of indefinite metrics*, Math. Ann., **232**(3), (1978), 267-272.

[12] M. Gutiérrez and B. Olea, *Global decomposition of a Lorentzian manifold as a generalized Robertson-Walker space*, Differential Geom. Appl., **27**(1), (2009), 146-156.

[13] M. Gutiérrez, F. J. Palomo and A. Romero, *A Berger-Green type inequality for compact Lorentzian manifolds*, Trans. Am. Math. Soc. **354**(11), (2002), 4505-4523.
[14] S. G. Harris, *A triangle comparison theorem for Lorentz manifolds*, Indiana Math Journal **31**(3), (1982), 289-308.
[15] S. G. Harris, *A characterization of Robertson-Walker spaces by null sectional curvature*, Gen. Relativity Gravitation, **17**(5), (1985), 493-498.
[16] S. G. Harris, *Some comparison theorems in the geometry of Lorentz manifolds*, Ph. D. Thesis, University of Chicago (1979).
[17] S. W. Hawking, S. W. and G. F. R. Ellis, *The Large Scale Structure of Space-time*, Cambridge University Press, UK, (1973).
[18] O. Kobayashi and M. Obata, *Certain mathematical problems on static models in general relativity*, Proceedings of the 1980 Beijing Symposium on Differential Geometry and Differential Equations, **3**, (eds. S. S. Chern and W. Wen-tsün), (1980), 1333-1344.
[19] L. Koch-Sen, *Infinitesimal null isotropy and Robertson-Walker metrics*, J. Math. Phys. **26**, (1985), 407-410.
[20] R. S. Kulkarni, *The values of sectional curvature in indefinite metrics*, Comment. Math. Helv **54**, (1979), 173-176.
[21] A. H. Lightstone, *Linear Algebra*, The Appleton-Century Mathematics Series, New York, 1969.
[22] K. Nomizu, *Remarks on sectional curvature of an indefinite metric*, Proc. Amer. Math. Soc., **89**(3), (1983), 473-476.
[23] B. O'Neill, *Semi-Riemannian Geometry With Applications to Relativity*, **103**, Academic Press, New York, (1983).
[24] F. J. Palomo, *The fibre bundle of degenerate tangent planes of a Lorentzian manifold and the smoothness of the null sectional curvature*, Diff. Geom. and Appl., **25**(6), (2007), 667-673.
[25] F. J. Palomo, Francisco and A. Romero, *Certain actual topics on modern Lorentzian geometry*, Handbook of differential geometry. Vol. II., Amsterdam, (2006), 513-546.
[26] A. Romero and M. Sánchez, *On the completeness of certain families of semi-Riemannian manifolds*, Geometriae Dedicata **53**, (1994)(1), 103-117.
[27] M. Sánchez, *Geodesic connectedness of semi-Riemannian manifolds*, Nonlinear Analysis **47**, (2001), 3085-3102.

(B. R. Yavuz) Department of Mathematics, Bilkent University, Bilkent, 06800 Ankara, Turkey
E-mail address: bengi@fen.bilkent.edu.tr

(B. Ünal) Department of Mathematics, Bilkent University, Bilkent, 06800 Ankara, Turkey
E-mail address: bulentunal@mail.com

(F. Dobarro) Instituto de Desarrollo Económico e Innovación, Universidad Nacional de Tierra del Fuego, Antártida e Islas del Atlánti-co Sur, Gobernador Paz 1410, 9410, Ushuaia, Tierra del Fuego, Argentina.
FCE, Universidad Austral, Mariano Acosta s/n y Ruta Nac. 8, Edificio Grado, B1629WWA, Buenos Aires, Argentina
E-mail address: fdobarro@untdf.edu.ar, fdob07@gmail.com